Sixth Annual
Raymond N. Castle
Student Research Conference

Department of Chemistry

University of South Florida

2008



Sixth Annual
Raymond N. Castle
Student Research Conference

Department of Chemistry

University of South Florida

2008



Table of Contents

Welcome	2
Raymond N. Castle Research Conference Committee	3
Table of Contents	4
Sponsors	5
Schedule of Events	6
Raymond N. Castle	7
Plenary Speaker – Dr. Vern Schramm	8
Dr. Dean F. Martin, special thanks	10
Session Schedule	11
Poster Titles	13
Morning Session Abstracts	18
Afternoon Session Abstracts	21
Undergraduate Poster Abstracts	26
Graduate Poster Abstracts	35
Acknowledgements	41

Sponsors

We would like to thank our sponsors.

For additional sponsors, please see the insert in this program

CLEAR SPRINGS LAND COMPANY, LLC
690 E. Davidson Street
Bartow, FL 33830
(863) 534-1292

GRAY'S COLLEGE BOOKSTORE
2836 E Fletcher Ave
Tampa, FL 33612-9420
(813) 971-2665

Thank you to Gray's bookstore for their generous donation of printing these booklets for the Sixth Annual Raymond N. Castle Conference.

McALISTER'S DELI
11402 N. 30th Street
Tampa FL 33612
(813) 341-1160

Plenary Speaker

Dr. Vern Schramm, Albert Einstein College of Medicine



Dr. Vern Schramm, a native of South Dakota, earned his bachelor's degree from South Dakota State College. He then earned a Master's degree from Harvard University and a Ph.D. in the mechanism of enzyme action from the Australian National University. Upon returning to the United States, Dr. Schramm served as a NSF-NRC Postdoctoral Fellow at the NASA Ames Research Center. He then joined the faculty of Temple University School of Medicine, where he began fundamental research designed to understand enzymatic transition state structure. In 1987, he moved to the Bronx to become Professor and Chair of Biochemistry at Einstein. He was appointed to the Ruth Merns Chair of Biochemistry in 1995. At Einstein, his research on transition state structure has developed to become a powerful method for drug design.

Dr. Schramm is world-renowned for his research into the “transition-state structure” of enzyme-catalyzed reactions -- the shapes that reacting molecules assume when enzymes catalyze chemical reactions. As he describes it, “By knowing the transition-state structure of enzyme-catalyzed reactions, we can design powerful inhibitors of enzymes to be used as drugs or antibiotics.” Two of the inhibitors designed by the Schramm laboratory have entered clinical trials. One of the antibiotics shows promise for treating leukemia that does not respond to other therapy. The second is in clinical trials for eventual application to autoimmune diseases. These include psoriasis, rheumatoid arthritis, multiple sclerosis, inflammatory bowel disorders, insulin-dependent diabetes and tissue transplant rejection. New agents are being explored for malaria, solid tumors, and as bacterial antibiotics.

Dr. Schramm has received numerous honors in recognition of his contributions for both research and teaching in biochemistry and enzymology. These include a ten-year Merit Award from the NIH for work on transition states, the Repligen Award from the American Chemical Society, election as a Fellow of the American Association for the Advancement of Science, the Rudi Lemberg Award from the Australian Academy of Science, the Election to the Davidoff Society at Einstein, the George A. Sowell Award for Excellence in Teaching from Temple University School of Medicine, the Harry Eagle Award for Outstanding Basic Science Teaching from the Albert Einstein College of Medicine, and most recently his election into the National Academy of Sciences. Dr. Schramm has served as the Chair of the Division of Biological Chemistry of the American Chemical Society and is an Associate Editor of the Journal of the American Chemical Society. He serves as a scientific advisor to national research resources and several biotechnology companies.

Dr. Dean F. Martin, Special Thanks

Dean F. Martin is Distinguished Service Professor and Director of the Institute for Environmental Studies at the University of

Session Schedule

Morning Session (CHE 100)
Session Chair: Emma Farrell

9:00 – 9:15 **Chunyan Wang**
"The in vitro and in vivo Evaluation of Implanted Glucose Sensors with Hydrogel Coatings"

9:15 – 9:30 **Mariangela Dametto**
"Computer simulations of the refolding of apomyoglobin from high temperature unfolded state"

9:30 – 9:45 **Jeremy Beau**
"*Drug Discovery from Floridian Mangrove Endophytes"*

9:45 – 10:00 **Laura Anderson**
"Design and Synthesis of Peptidomimetics for the Disruption of Bcl-2-family Proteins"

10:00 –10:30 Break

10:30 – 10:45

Afternoon Session (CHE 100)
Session Chair: Jaime Heimbegner

2:45 – 3:00 **Sung Wook Yi**
"Cysteine based PNA (CPNA): Design, synthesis and Application"

3:00 – 3:15 **Ryan Cormier**
"Polyacrylate Nanoparticles for Potential Delivery of Chloroquine Resistance Reversal Agents"

3:15 – 3:30 **Ruizhi Wu**
"Novel synthesis and polymerization of protected dihydroxy 7-member cyclic carbonate"

3:30 – 3:45 **Divya Ramamoorthy**
"Inhibitors of SHP2, a protein tyrosine phosphatase"

3:45 – 4:00 **Matt Lebar**
"Degradation and Partial Reconstruction (C3-14) of Cytotoxic Macrolide Palmerolide A"

4:00 – 4:15 Break

4:15 – 4:30 **Arun Babu Kumar**
"Development of a Labeling Probe for the Discovery and Identification of Saccharide-Binding Proteins"

4:30 – 4:45 **Shawn Hamm**
"Role of the Disulfide Bridge on the Stabilization and Dynamics of Human Cu, Zn Superoxide Dismutase Upon Loss of Zn"

4:45 – 5:00 **Pasha Khan**
"Pd catalyzed allylic alkylation- Application towards synthesis of new bioactive compounds"

5:00 – 5:15 **Sameer Kulkarni**
"Application of Target Guided Synthesis (TGS) approach : Targeting protein-protein interactions"

5:30 Awards Ceremony

17. **Nicholas Kondis**
    "Removal of aqueous nuisance ions using Cuprilig(copper attached to immobilized ligands, IMLIGs)"
18. **Jason Kuczynski**
    "Using Alkoxy Bridged Tetracarboxylates to Create Novel MOFs"
19. **Michael Manasterski**
    "

The Barbara and Dean F. Martin Graduate Poster Session

(NES First Floor)

34. **Jonathan Belof**
"Canonical Monte Carlo simulation of a Metal-Organic Framework utilizing a polarizable potential"
35. **Ryan Cormier**
"Polyacrylate Nanoparticles for Potential Delivery of Chloroquine Resistance Reversal Agents"
36. **Michelle Cortes-Salva**
"The Design, Synthesis and Evaluation of 1,3-di-o-tolylguanidine (DTG) Analogues as Potential Active Anti-Stroke Therapeutics"
37. **Emma Farrell**
"Uncovering the biosynthetic pathway of primary fatty acid amides"
38. **Shawn Hamm**
"The Role of the Disulfide Bridge on the Conformational Relaxation of Cu, Zn Superoxide Dismutase Upon Loss of Metals Studied by Molecular Dynamics"
39. **Milena Ivkovic**
"N-Acylethanolamines as substrates for ADH3"
40. **Pasha M Khan**
"Pd(0)-catalyzed intramolecular alkylation: Stereoselective synthesis of furan and isoxazoline-2-oxide analogs
41. **Matt Lebar**
"On the stereochemistry of palmerolide A"
42. **Edward W. Lowe, Jr.**
"Elucidation of ascorbate binding sites in peptidylglycine α-amidating monooxygenase through molecular dynamics simulations"
43. **Shikha Mahajan**
"Synthesis of biotinylated-azido-Adenine-Ribose derivatives analogues: Potential activity based protein profiling probes"
44. **John Maschek**
"Marine extracts with antiviral activity"
45. **David Z. Myers**
"Design and synthesis of piperazines with potential modulation of Bcl-X_L-protein interactions"
46. **William A. Maza**
"Photophysical properties of three alyx[4]resorcinarenes of amphiphilic character"
47. **John Perry**
"Building with Bigger Blocks: Nanoscale molecular faceted polyhedra as nodes in extended metal-organic materials"
48. **Joshua V. Ruppel**
"Cobalt-Catalyzed Asymmetric Cyclopropanation"

49. Sung Wook Yi

"Regioselective Mono Acylation of the Electronically Less Reactive Nitrogen of

Session Abstracts

Saturday, April 19th, 2008

Morning Session (CHE 100)

Session Chair: Emma Farrell

9:00 – 9:15 **Chunyan Wang**

"The in vitro and in vivo Evaluation of Implanted Glucose Sensors with Hydrogel Coatings"

In order to protect implanted glucose sensors from biofouling, novel hydrogels (VP30) produced from a monomeric mixture of 34.5% HEMA, 34.5% DHPMA, 30% VP and 1% EDGMA were coated on two kinds of VP30-coated sensors, Pt/GOx/VP30 and Pt/GOx/Epoxy-Polyurethane (EPU)/VP30 sensors. They were examined in glucose solutions during a period of 4 weeks. The hydrogel coating was about 40 μm thick and was firmly attached to both types of sensors during experiments. Due to the poor response linearity of the Pt/GOx/VP30 sensors, the hydrogel coating could not be used as a diffusion-limiting layer. Therefore, the further studies were focused on the Pt/GOx/EPU/VP30 sensors. With a diffusion-limiting epoxy-polyurethane membrane, the linearity was improved (2-30 mM) and the response time was within 5 min. Eight Pt/GOx/EPU/VP30 sensors were subcutaneously implanted in rats and tested once per week over 4 weeks. 100% of the implanted sensors kept functioning for at least 21 days and 3 out of 8 sensors still functioned at day 28. We previously tested 68 Pt/GOx/EPU sensors in rats and only 13.5% of the sensors survived at day 21. The histology analysis revealed that the fibrous capsule formed surrounding the hydrogel-coated sensor was thinner than that formed around the Pt/GOx/EPU sensor. Inflammatory cells were no longer present in the capsular tissue when the sensor was coated with hydrogels.

9:15 – 9:30 **Mariangela Dametto**

"Computer simulations of the refolding of apomyoglobin from high temperature unfolded state"

Folding mechanisms of apomyoglobin were studied by performing computer simulations with atomic detail. The mechanisms of folding of this protein have been extensively

9:30 – 9:45 **Jeremy Beau**

"*Drug Discovery from Floridian Mangrove Endophytes"*

Recent studies have shown that within the leaves, bark, roots and seeds of mangroves exists an immense world of endophytic organisms. These microscopic communities are complex, providing a great diversity of secondary metabolites with special functions that combat the microbe-plentiful seawater. These secondary metabolites have the potential to be potent and highly selective drug candidates. A collection of endophytes from Floridian mangroves is currently being developed. The microorganisms are cultured in the laboratory and then systematically screened against various illnesses such as cancers, infectious diseases and common pathogenic microbes. After novel compounds are identified they can be characterized by NMR or X-Ray diffraction. Finally, synthetic routes from commercially available reagents can be explored. This paper reports on the preliminary collection of these organisms and the results obtained from the bioassays.

9:45 – 10:00 **Laura Anderson**

"Design and Synthesis of Peptidomimetics for the Disruption of Bcl-2-family Proteins"

Disregulation of the apoptotic processes is one of the major causes of diseases, such as cancer, inflammatory, autoimmunity, and neurodegenerative disorders. Within the B-cell lymphoma-2 (Bcl-2) family, the balanced interactions between anti and pro-apoptotic proteins play a major role in regulating apoptosis. Several studies have indicated that overexpression of anti-apoptotic Bcl-2 and Bcl-xL proteins is associated with tumor progression and drug resistance. Previously reported compounds that inhibit the interaction between pro-apoptotic BH3 domains and Bcl-2 family proteins contain hydrophobic scaffolds that can diminish their use as potential drug leads. Herein, we present the design and synthesis of hydrophilic alpha-helix mimics based on hydrazine linked piperazine-dione repeat units for the disruption of Bcl-2 proteins. Several of these compounds have been synthesized in4 TDar[zed ifl( )]TJ resis ha-ic

11:00 – 11:15 **Christi Young**

"Molecular Dynamics Simulations of Influenza A Non-structural Protein 1 (NS1) RNA Binding Domain"

Six 50-ns molecular dynamics simulations of the RNA binding domain of the Non-structural protein 1 (NS1) of influenza A virus, a homodimer, were performed at 298K. We focused our analysis on helices 2 and 2', which are involved in RNA binding. A salt

Afternoon Session (CHE 100)

Session Chair: Jaime Heimbegner

2:45 – 3:00 **Sung Wook Yi**

"Cysteine based PNA (CPNA): Design, synthesis and Application"

Peptide nucleic acids (PNA), a pseudopeptide DNA mimic, was discovered by Nielsen and his coworker in 1991. PNA is proved to sequence-specifically form a very stable duplex with complementary DNA and RNA strands through Watson-Crick base paring, and it is also capable of binding to duplex DNA by helix invasion. However, due to its acyclic, achiral and neutral nature of the backbone, PNA has shown problems such as its poor aqueous solubility, poor cell permeability and instability of PNA-DNA duplexes and triplexes. Accordingly, many synthetic approaches have been directed toward developing modified backbones of PNA. Among those PNA analogs, only few examples including lysine-based monomers, guanidine-based peptide nucleic acids (GPNA) and the aminoethylprolyl PNA (aep-PNA) showed noticeable enhancements with regards to the daunting challenges mentioned above. I will present our research endeavor to develop the CPNA oligomers with the great water-solubility and cell permeability.

3:00-3:15 **Ryan Cormier**

3:30 – 3:45 **Divya Ramamoorthy**
"Inhibitors of SHP2, a protein tyrosine phosphatase"

Tyrosine phosphatases play an important role in regulating the phosphorylation statu:l iu04(a)3i1c

4:15 – 4:30 **Arun Babu Kumar**

"Development of a Labeling Probe for the Discovery and Identification of Saccharide-Binding Proteins"

Proteins that interact with carbohydrates in a non-covalent fashion occur widely in nature. Such proteins, which have come into the forefront of biological research in recent years, belong to the class of the lectins. We discuss the design and development of a

4:45 – 5:00 **Pasha Khan**

“Pd catalyzed allylic alkylation- Application towards synthesis of new bioactive compounds”

Pd catalyzed alkylation of allylic substrates is one of the most synthetically useful reactions and has been extensively used in organic synthesis. Hence, in quest of synthesizing new bioactive compounds, organopalladium chemistry was utilized in conjunction with enzyme catalyzed reactions. New optically pure isoxazoline-2-oxide and furan analogs can be synthesized starting from a meso-diol via Pd(0) catalysis. Pd(0) catalyzed allylic alkylation was also applied to nitroalkanes and other nitroesters to obtain unnatural amino acids and amino alcohols. This methodology is compatible with wide spectrum of functional groups like NMslogy i(. New)

Poster Exhibit Abstracts

production of certain types of polymers. It is necessary to know the magnitude of the charge an

12. Kenneth Grandalski

"Neutral and Anionic Metal-Organic Frameworks from Co-crystal Controlled Solid State Ligands"

Co-crystal Controlled Solids State Synthesis (C3S3) reactions produce cyclic imides in high yields while reducing or eliminating solvents and/or solvent waste. Benzoimide Phenanthroline m-Dicarboxylic Acid (H2BIPA-DC) was choosen for this synthesis because it possesses similarities to other dicarboxylic acids currently used for Metal-Organic Frameworks (MOF's). The H2BIPA-DC from solvent-drop grind of 1,4,5,8-Napthelentetracarboxylic Dianhydride and 3-Aminobenzoic

replacement of the TZI bridging ligand and TZI-copper trimer with 5,5',5''-[1,3,5-phenyltri(methoxy)]tri-isophthalic acid (H6BTMOI). Comparisons between these two iso-structures will be made in reference to crystal synthesis, crystal porosity, and hydrogen adsorption properties.

16. Benjamin Holt

"Effect of High Temperature on the Structure of Cu, Zn Superoxide Dismutase Studied with Molecular Dynamics"

In this study we simulated the essential human antioxidant enzyme, Cn, Zn superoxide dismutase. Immature and mutant forms of the enzyme may misfold and aggregate. These aggregates are believed to be toxic, and are linked to Familial Amyotrophic Lateral Sclerosis. Four 5-ns Isobaric-Isothermal ensemble molecular dynamics trajectories of the wild type apo protein were created at 300K, 305K, 310K, and 315K. Fluctuations in the structure of the enzyme were studied as a function of temperature.

17. Nicholas Kondis

"Removal of aqueous nuisance ions using Cuprilig(copper attached to immobilized ligands, IMLIGs)"

The present study describes the synthesis of a copper(II) salt of a commercially available immobilized ligand (IMLIG), Octolig-21® and the efficacy of this composite, called Cuprilig to remove various nuisance ions from aqueous solutions. The ions attempted in this study include nitrate, nitrite, perchlorate, nitrogen (as ammonium chloride), sulfate, arsenate, chromate, and phosphate. The synthesis was accomplished by treating an aqueous suspension 82 g of Octolig-21® in 150 ml of deionized water with 0.2 moles of copper(II)acetate monohydrate in 450 ml, shaken overnight, filtered through a sieve, and rinsed to remove excess copper salts. Standard solutions of the aforementioned ions were prepared in various concentrations ranging from 30 ppm

23. Hai Nguyen

"The effect of mutation of residues on molecular bynamics of the RNA binding domain of influenza A virus Non structural protien 1"

The main goal of this project is to study the effect of mutations of specific residues on the interactions between the monomers of the RNA binding domain of the influenza A virus non structural protein 1, a homodimer. Several specific Alanine mutations,such as Arginine 46, Arginine 38, and Aspartate 29, were simulated by computer programs. The molecular dynamics simulations were performed through several steps; were minimization, heat, equilibriation, and then run. Each step required certain input files in order to run. After the simulations finished, the distances between atoms of residues involved in the mutationswere determined through the VMD program. The distances measured were compared to distances from the original, non-mutated RNA binding domain.

24. Patrick Nugent

"Co-crystal Controlled Solid-State Synthesis of a Diimide, Designed for Supramolecular Chemistry"

Benzoimide pyromellitic tetracarboxylic acid, H4BIPRO-TC, has been formed via co-crystal controlled solid-state synthesis, C3S3, for the purpose of creating a species that can readily form co-crystals and coordination polymers. This diimide synthesis is carried out by grinding reactants in minute amounts of solvent, culminating in high yields while creating little to no waste. The product was characterized via IR, PXRD, 1H-NMR, and x-ray single crystal diffraction. H4BIPRO-TC was functionalized with carboxylic acids to participate in non-covalent interactions (i.e. hydrogen bonding) that have been shown to sustain supramolecular synthons. The species was observed to form a 1:2 cocrystal with 4,4'-dipyridyl. By design, H4BIPRO-TC has a sterically limited torsional angle between the imide and N-phenyl planes upon crystallization.

25. Amanda Okolo

"Bio-catalytic Transformations in Organic Synthesis- A route to new bioactive compounds "

The proposed work involves the synthesis of new, optically pure compounds through the use of lipases (hydrolytic enzymes). This work further enhances the importance of green chemistry in organic synthesis. The synthesis will involve the use of commercially available cyclic compounds, which will be converted to meso diacetates in high yield. These diacetates will be hydrolyzed using hydrolytic enzymes namely lipases to yield mono esters in high yield and enantioselectivity. The final enantiopure products will serve as starting material for synthesis of new bioactive compounds.The main emphasis of this work will be to limit the use of expensive metal catalysts and induce high enantioselectivity via biocatalysis, thereby promoting the concept of green chemistry.

26. Nik-Chay Reithma

"Germania Based Capillary Microextraction (CME)"

Traditional gas and liquid chromatography use silica based stationary phases for extraction. The traditional stationary phases can pose problems when looking at particularly polar analytes or for

focused on extending the family f germania based stationary phases and developing germania based monoliths using PEG-600 and TMOS.

27. Ian-Anthony Rusiana

"Design and Synthesis of Cyclic β-Hairpin Peptides as MDM2-P53 Interaction Inhibitors"

Many proteins exhibit protein-protein interactions involving relatively small regions of thei

light microscopy, and species identification by polymerase chain reaction analysis. Preliminary

Graduate Posters (NES First Floor)

obtain.


Pd(0)
CO_2C_2
N
O


PHM
PAL
O
R
N
H
COOH
O_2
H_2O
2 Cu(II)
2 Ascorbate
2 Semidehydroascorbate
OH
Glyoxylate
Zn(II)
NH_2

45. William A. Maza

"Photophysical properties of three calix[4]resorcinarenes of amphiphilic character"

Here we report the photophysical properties of three resorcinol derived cavitands. Steady-state absorption and fluorescence spectra for the three cavitands studied displayed absorption maxima ranging from 270nm to 290nm and emission maxima between 300nm and 320nm. The absorption/emission spectra are attributed to excitation of the substituted phenyl rings that constitute the cavitand cavity. The similarity of the absorption/emission spectra and fluorescence quantum yields between the cavitands and 5-hydroxy-1,3-benzenedicarboxylic acid (Hbdc), an analoguos substituted benzene, indicate weak intramolecular interactions between the monomers of the cavitand core. Fluorescence lifetimes were found to increase from ~4 ns to ~12 ns upon functionalization with groups with greater electron donating ability. Combined with polarization data, we find that the inclusion volume is related to the O···H interactions at the upper rim of the cavitand.

46. David Z. Myers

"Design and synthesis of piperazines with potential modulation of Bcl-X_L-protein interactions"

001lnreg 7aaorspop. 8.9285(o)-5(n w)-4pipy h4 >>v(12)b288 valid the a(-)tw -t the v(1targe)-5forhe cavitar

48. Joshua V. Ruppel

"Cobalt-Catalyzed Asymmetric Cyclopropanation"

Metal-catalyzed cyclopropanation of olefins with diazo reagents has attracted great research interest because of its fundamental and practical importance. The resulting cyclopropyl units are recurrent motifs in biologically important molecules and can serve as versatile precursors in organic synthesis. We have previously reported the cobalt-catalyzed cyclopropanation of styrene and styrene derivatives with diazo reagents. With the use of a new class of D_2-symmetric chiral porphyrin ligands, the resulting cyclopropane units are produced in high yields with excellent diastereo- and enantioselectivities. The dimerization of diazo compounds is minimized in the cobalt-mediated system, allowing for a one-pot protocol and eliminating the need for slow addition of diazo reagents. In our continuing effort to further improve and expand the scope of the cobalt-catalyzed system, we will present our new results on asymmetric cyclopropanation of electron-deficient olefins and the use of diazo reagents other than ethyl diazo acetate and *t*-butyl diazo acetate.

49. Sung Wook Yi

"Regioselective Mono Acylation of the Electronically Less Reactive Nitrogen of Aryl Hydrazines via Temporary Protection with Carbon Disulfide"

Hydrazines are an important class of compounds that contain the reactive nitrogen-nitrogen bond. Hydrazines and analogues such as, hydrazides and hydrazones, are well known for their highly active biological profiles. For example, a substantial number of hydrazines and their analogues are known to have antineoplastic activity. Hydrazines are also frequently used in synthetic chemistry to make pyrazoles, oxadiazoles, triazoles and indoles. However, alkylating and acylating hydrazines can be cumbersome especially when onE29 0 T54[(Hydrazines and aall)1ono-a

Acknowledgments

Dr. Jon Antilla

Dr. Bill Baker

Dr. Kirpal Bisht

Dr. Wayne Guida

Dr. Julie Harmon

Dr. Milton Johnston

Dr. Vicky Lykourinou

Dr. Roman Manetsch

Dr. David Merkler

Dr. Patricia Muisener

Dr. Peter Zhang

University of South Florida
Department of Chemistry